

Cytotoxicity Studies of *Mortierella alpina* AIB RR2 Extracts on HeLa and A549 Cell Lines

Arunchand Rayaroth, Rajesh Singh Tomar, Raghavendra Kumar Mishra

Department of ??? Amity Institute of Biotechnology, Amity University Madhya Pradesh, Maharajpura (Opposite Airport), Gwalior, Madhya Pradesh, India

AQ1

Abstract

Aim: The aim is to understand any possible cytotoxic activity of *Mortierella alpina* extract against carcinoma cell lines and possible interaction with normal cell lines. **Materials and Methods:** The extract of *M. alpina* at varying concentrations was incubated with the normal Vero cell lines and also the carcinoma cell lines HeLa and A549, after which 3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyl tetrazolium bromide assay was carried out to determine the viable cell. **Results and Discussion:** The cytotoxic activities of *M. alpina* extract were quite strong against both HeLa and A549 carcinoma cell lines with inhibitory concentration 50 values of 106 and 116 µg/mL, respectively. However, the cytotoxic concentration 50 value of the fungal extract against the normal Vero cell line was 2674 µg/mL, which shows the very low toxic activity of the fungal extract against the normal cell lines. Further, the therapeutic index of *M. alpina* against HeLa and A549 cell lines was found to be 25.22 and 23.02, respectively. **Conclusion:** The present investigation indicates the probable use of *M. alpina* extracts in the treatment of carcinoma.

Key words: A549, HeLa, *Mortierella alpina*, 3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyl tetrazolium bromide assay, therapeutic index

INTRODUCTION

Mortierella alpina, a filamentous fungus generally found in saprophytic soil and has oleagenic properties, is a widely distributed psychrophilic Mortierellae.^[1-3] This fungus with a rosette globe pattern under cultural conditions is used extensively to produce omega fatty acids mostly under submerged fermentation. *M. alpina* is considered to have a promising future in the development of production related to omega fatty acid, especially in the omega 6 group.^[4] Due to the presence of important secondary metabolites, this oleaginous microorganism (psychrophilic filamentous) is presently highly considered for large-scale production of arachidonic acid (ARA).^[4,5]

Microbial metabolites have a long history of usage as therapeutic agents. Microbial metabolites used as therapeutic agents include antibiotics, enzymes, cholesterol-lowering metabolites, anticancer agents, and immunosuppressants.^[6] Although the number of potent and active metabolites discovery has

seen a growing trend, the requirement for novel metabolites is growing at a faster pace due to the increasing trend in untreatable diseases including cancer. Further, any major discoveries have to be confirmed not only for its efficacy but also for its safety before introducing it into a treatment regime. This is because of the fact that the active compound as well as any minor metabolite associated with the fungus might have some toxic effects on the normal functioning of cell and tissues.^[7] Preliminary safety studies are generally conducted on cell lines isolated from healthy mammalian tissues, whereas the effectiveness and potential of the active component to treat various diseased conditions are tested on specific affected cell lines including carcinoma cells.^[8,9]

According to the WHO, 8.8 million deaths reported in 2015 worldwide due to different types of cancer.^[10] Chemotherapy

Address for correspondence:

Raghavendra Kumar Mishra, Amity Institute of Biotechnology, Amity University Madhya Pradesh, Gwalior. Phone: +91-7415444342. E-mail: ???

AQ2

Received: 18-08-2018

Revised: 28-08-2018

Accepted: 12-09-2018

is one of the few potential and effective treatment strategies available.^[11] **With effectiveness mainly restricted to the preliminary stages.** Functional foods capable of supporting health and preventing and protecting the biological system^[12] from fatal diseases such as cancer are the need of the hour. *M. alpina*, which has been considered as a gifted microorganism to produce the conditionally essential fatty acid ARA, with numerous role in the physiological system including development and prevention of age-related brain and cardiovascular development as well as cognitive functions,^[4,13] still more need to be exploited for its numerous other potential roles in promoting health and preventing diseases. Due to the proven role of ARA in improving and preventing various degenerative diseases and also due to a long history of fungal metabolites in preventing cancer and related diseases, *M. alpina* can be a potent candidate to be evaluated for its anticancer properties. The present study is focussed on the cytotoxic and anticancer properties of a fungal isolate *M. alpina* AIB-RR2.

MATERIALS AND METHODS

All the media chemicals including Potato Dextrose Agar, Potato Dextrose Broth (PDB), glucose, and polysorbate 80 were purchased from HiMedia (Mumbai, India). Cell lines HeLa, A459, and Vero cell lines were purchased from the National Center for Cell Science, Pune, cell culture medium, Dulbecco's Modified Eagle's Medium - high glucose (AL111, HiMedia), Adjustable Multichannel Pipettes and a Pipettor (Eppendorf, Germany), fetal bovine serum (RM10432, HiMedia), 3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyl tetrazolium bromide (MTT) reagent (5 mg/mL) (4060 HiMedia), dimethyl sulfoxide (DMSO) (PHR1309, Sigma), Camptothecin (C9911, Sigma), D-PBS (TL1006, HiMedia), 96-well plate for culturing the cells (From Corning, USA), T25 flask (12556009, Biolite - Thermo), 50 mL centrifuge tubes (546043 TARSON), 1.5 mL centrifuge tubes (TARSON), 10 mL serological pipettes (TARSON), 10–1000 μ L tips (TARSON), 96-well enzyme-linked immunosorbent assay (ELISA) plate reader or spectrophotometer capable of measuring the absorbance (ELX-800 Biotek), inverted microscope (Biolink), and 37° C incubator with humidified atmosphere of 5% CO₂ (Heal Force, China) were the reagents and materials used in the present study.

Preparation of fungal extract

M. alpina AIB RR2, obtained from Mysore, Karnataka, and characterized using internal transcribed spacer sequencing, was grown on a PDB medium for 7 days at 25° C. On completion of incubation, mycelia were harvested and mixed with ethyl acetate and kept for vigorous shaking for 1 h. On completion of extraction, the supernatant was collected and concentrated under vacuum using a rotary evaporator at 40°C and dried under nitrogen. The concentrate was stored at –80°C for future experimentation.^[14]

Cytotoxicity analysis of the selected strain by MTT assay

About 200 μ L of the culture of suspension cell (cell density 20,000 cells per well) was seeded into a 96-well plate (ELISA plate) and was allowed to grow for about 12 h. After completion of incubation, appropriate concentrations of the test agents ranging from 0.1 to 1000 μ L were added to the plates and incubated for 24 h at 37°C in a 5% CO₂ atmosphere. After completion of the incubation period, the ELISA plates were removed from the CO₂ incubator and spent media were washed with D-PBS. MTT reagent was added to this to a final concentration of 0.5 mg/mL of total volume, and the plate was wrapped with aluminum foil to avoid exposure to light and incubated for 3 h. On completion of incubation, the excess MTT reagent was washed and then added 100 μ L of DMSO as a solubilizing agent with intermittent gentle stirring in a gyratory shaker. The controls included were medium control (medium without cells), negative control (medium with cells but without the experimental extract), and positive control (medium with cells and with 50 μ M of CPT/75 μ M of H₂O₂). The absorbance was read on an ELISA reader at 570 nm with 630 nm used as reference wavelength. The percentage inhibition was calculated using the following equation:

$$\% \text{ Cell inhibition} = \frac{(\text{Absorbance value of control} - \text{Absorbance value of sample})}{\text{Absorbance value of control}} \times 100$$

The inhibitory concentration (IC₅₀) value which denotes the inhibition of 50.0% of the cancer cell growth and cytotoxic concentration (CC₅₀) which denotes the inhibition of 50.0% of the healthy cell growth was determined using linear regression equation, i.e. $y = mx + b$, with $y = 50$, and m , x and b values were derived from the viability graph. The therapeutic index (TI) value was derived^[15,16] by the following equation:

$$TI = \frac{CC_{50}}{IC_{50}}$$

RESULTS

In the present study, the *in vitro* cytotoxic effects of *M. alpina* AIB-RR2 extracts on HeLa and A549 cells were assessed. The cytotoxicity of the fungal extracts was also investigated against the normal Vero cell line. It was found that *M. alpina* AIB-RR2 exerts a concentration-dependent antiproliferative effect on the tested carcinoma cell lines. The extract showed an exponential increase in cytotoxic activity against both HeLa and A549 cells. The IC₅₀ value of the extract against HeLa cells was 106 μ g/mL. A similar result was also found against A549 cells, which showed the IC₅₀ value of 116 μ g/mL. In both the cytotoxic activity graphs, slope showed a very high correlation between increasing concentration of the fungal extract and corresponding decrease in cell viability as depicted by the R² value of 0.977–0.971 with HeLa and A549 cells, respectively.




Figure 1: Concentration versus % viability with Vero cells




Figure 4: Microscopy of cytotoxicity analysis in A549 cells using dimethyl sulfoxide extract of *Mortierella alpina* (a) at 1.0 µg and (b) at 125 µg




Figure 2: Concentration versus % viability with HeLa cells




Figure 5: Microscopy of cytotoxicity analysis in HeLa cells using dimethyl sulfoxide extract of *Mortierella alpina* (a) at 1.0 µg and (b) at 125 µg (*Scale Bar = 100 µm)




Figure 3: Concentration versus % viability with A459 cells

Table 1: Cytotoxicity analysis in HeLa cells using DMSO extract of *M. alpina*

Cell line	IC ₅₀ values (in µg/mL)	CC ₅₀ value (in µg/mL)	TI
HeLa	106		25.22
A549	116		23.05
Vero		2674	16.901

M. alpina: *Mortierella alpina*, DMSO: Dimethyl sulfoxide, IC: Inhibitory concentration, CC: Cytotoxic concentration, TI: Therapeutic index

Further, the non-inhibitory activity of the extract against the healthy Vero cell lines was very evident with a CC₅₀ of 2674 µg/mL. Furthermore, the TI value which is the ratio of CC₅₀ and IC₅₀ was recorded at 25.22 and 23.05 with HeLa and A549 cells, respectively.

Cytotoxicity data

???

DISCUSSION

The increasing number of cancer cases around the world is demanding for more improved strategy for prevention

as well as treatment of this dreaded disease. Approaches, which are more than just a treatment mode rather which may be included as a nutritional supplement in our regular diet, can be a better strategy to prevent carcinogenesis. Fungal extracts including *Saccharomyces cerevisiae* and the endophytic fungus *Annona muricata* extracts have been demonstrated previously to possess anticancer activity in cancer cell lines *in vitro*.^[14,17] The present study reports the anticancer activity of the oleaginous fungus, *M. alpina*, which is also having one of the most efficient ARA-producing capabilities. The extracts of *M. alpina* which has been previously reported to have multiple biological activity including antioxidant and antibacterial activity^[18] showed a good inhibitory effect on the cervical cancer cell line HeLa, as well as on the adenocarcinoma cell line A549. Further, the low IC₅₀ of the extract against both the cell lines highlights the potential of the extracts as a possible therapeutic agent. The IC₅₀ values obtained in the present study were in line with the maximum limit set for crude extracts at 30 mg/mL by the National Cancer

Institute.^[19] This was ascertained by *in vitro* study carried out on the normal kidney Vero cell lines in which the extracts showed CC only at a very high concentration which was quite promising. Furthermore, the TI value reaffirmed the possible therapeutic ability of *M. alpina* AIB-RR2 extracts. As a result, it is predicted that this extract may possess significant therapeutic characteristics for cancer. At relatively low concentrations, this extract was able to inhibit HeLa cells and A549 cells and this inhibition was in a dose-dependent manner. This important finding gives a hope that *M. alpina* RRB₂ active materials can be useful in cancer. Further detailed studies needed to be conducted on the extracts of *M. alpina* to identify the potent anticancer molecules so as to develop specific extraction procedures as well as biotechnological approaches to improve the production of the target metabolite. Fungus produces numerous secondary metabolites, many of which are found to have multiple biological activities, which includes anticancer activities also. Some of the anticancer compounds produced by fungus include anguidine and rhizoxin which act at molecular level including protein synthesis inhibition and inhibition of microtubule assembly. Referring to the above-detailed anticancer secondary metabolites produced by filamentous fungi including *Fusarium* and *Rhizopus* spp., further detailed investigation needs to be conducted at the molecular level, including secondary metabolite separation and structural characterization to identify the target molecule. Further cellular level studies also need to be carried out to identify the specific possible mechanisms of action of the anticancer molecule produced by *M. alpina*, which might include inhibition of protein synthesis, inhibition of DNA synthesizing enzymes, and inhibition of angiogenesis. Furthermore, rigorous biotechnological studies have to be undertaken to identify and explore specific production, extraction, and separation procedures to improve the production of the target metabolite.

ACKNOWLEDGMENT

We acknowledge Dr. Ashok Kumar Chauhan, President, Redondo Beach Educational Foundation (RBEF), parent organization of Amity University Madhya Pradesh (AUMP), Dr. Aseem Chauhan, Additional President, RBEF, and chairman of Amity University Gwalior Campus, and Lt. Gen. V.K. Sharma, AVSM (Retd.), Vice-Chancellor of AUMP Gwalior Campus, for providing necessary facilities their valuable support and encouragement throughout the work.

REFERENCES

- Wagner L, Stielow B, Hoffmann K, Petkovits T, Papp T, Vágvolgyi C, *et al.* A comprehensive molecular phylogeny of the *mortierellales* (Mortierellomycotina) based on nuclear ribosomal DNA. *Persoonia* 2013;30:77-93.
- Wua T, Chellemia DO, Martinc KJ, Grahamb JH, Rosskop EN. Discriminating the effects of agricultural land management practices on soil fungal communities. *Soil Biol Biochem* 2007;39:1139-55.
- Samadlouie HR, Esfahani ZH, Alavi SM, Najafabadi MS, Sahari MA, Abbasi S. Statistical approach to optimization of fermentative production of oil and arachidonic acid from *Mortierella alpina* CBS 754.68. *Afr J Microbiol Res* 2012;6:1559-67.
- Rayaroth A, Tomar RS, Mishra RK. Arachidonic acid synthesis in *Mortierella alpina*: Origin, evolution and advancements. *Proc Natl Acad Sci India B Biol Sci* 2017;87:1053.
- Chen R, Matsui K, Ogawa M, Oe M, Ochiai M, Kawashima H, *et al.* Expression of $\Delta 6$, $\Delta 5$ desaturase and GLELO elongase genes from *Mortierella alpina* for production of arachidonic acid in soybean [*Glycine max* (L.) Merrill] seeds. *Plant Sci* 2006;170:399-406.
- Bérdy J. Bioactive microbial metabolites. *J Antibiot (Tokyo)* 2005;58:1-26.
- Martinez KB, Leone V, Chang EB. Microbial metabolites in health and disease: Navigating the unknown in search of function. *J Biol Chem* 2017;292:8553-9.
- Sodde VK, Lobo R, Kumar N, Maheshwari R, Shreedhara CS. Cytotoxic activity of *Macrosolen parasiticus* (L.) danser on the growth of breast cancer cell line (MCF-7). *Pharmacogn Mag* 2015;11:S156-60.
- Sadeghi-Aliabadi H, Minaiyan M, Dabestan A. Cytotoxic evaluation of doxorubicin in combination with simvastatin against human cancer cells. *Res Pharm Sci* 2010;5:127-33.
- World Health Organization. International Agency for Research on Cancer [IARC]. Latest World Cancer Statistics. Geneva, Switzerland: World Health Organization; 2013.
- Park IH, Lee KS, Ro J. Effects of second and subsequent lines of chemotherapy for metastatic breast cancer. *Clin Breast Cancer* 2015;15:e55-62.
- Aghajanpour M, Nazer MR, Obeidavi Z, Akbari M, Ezati P, Kor NM, *et al.* Functional foods and their role in cancer prevention and health promotion: A comprehensive review. *Am J Cancer Res* 2017;7:740-69.
- Kiso Y. Pharmacology in health foods: Effects of arachidonic acid and docosahexaenoic acid on the age-related decline in brain and cardiovascular system function. *J Pharmacol Sci* 2011;115:471-5.
- Minarni, Artika IM, Julistiono H, Bermawie N, Riyanti EI, Hasim, *et al.* Anticancer activity test of ethyl acetate extract of endophytic fungi isolated from soursop leaf (*Annona muricata* L.). *Asian Pac J Trop Med* 2017;10:566-71.
- Lin L, Hwang PL. Antiproliferative effects of oxygenated sterols: Positive correlation with binding affinities for the antiestrogen-binding sites. *Biochim Biophys Acta* 1991;1082:177-84.
- Hasan AE, Manguwidjaja D, Sunarti TC, Suparno O, Setiyono A. Investigating the antioxidant and

AQ5
AQ5

- anticytotoxic activities of propolis collected from five regions of Indonesia and their abilities to induce apoptosis. Emir J Food Agric 2014;26:390-8.
17. Senthilraja P, Kathiresan K. *In vitro* cytotoxicity MTT assay in Vero, HepG2 and MCF-7 cell lines study of marine yeast. J App Pharm Sci 2015;5:80-4.
 18. Melo IS, Santos SN, Rosa LH, Parma MM, Silva LJ, Queiroz SC, *et al.* Isolation and biological activities of an endophytic *Mortierella alpina* strain from the Antarctic moss *Schistidium antarctici*. Extremophiles 2014;18:15-23.
 19. Suffness M, Pezzuto JM. Assays related to cancer drug discovery. In: Hostettmann K, editor. Methods in Plant Biochemistry: Assays for Bioactivity. Vol. 6. London: Academic Press; 1990. p. 71-133.

Source of Support: Nil. **Conflict of Interest:** None declared.

Author Queries???

AQ1:Kindly provide department

AQ2:Kindly provide corresponding email id

AQ3:Kindly review the sentence as it seems to be incomplete.

AQ4:Kindly cite figures 1-5 in the text part

AQ5:Kindly provide author initial

AQ6:Kindly cite table1 in the text part

AQ7:Kindly provide the text part